



When depression sets you back, *Keep Moving Forward.*

Now that you've talked to your doctor, here's an introduction to adding REXULTI.



For eligible commercially-insured patients

THE REXULTI SAVINGS CARD:
Benefits for REXULTI
and your generic
ANTIDEPRESSANT.*

*Separate benefits. Terms and Conditions apply.

INDICATION: REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

SELECT IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis without agitation that may happen with dementia due to Alzheimer's disease.
- Increased risk of suicidal thoughts and actions. REXULTI and antidepressant medicines may increase suicidal thoughts and actions in some people 24 years of age and younger, especially within the first few months of treatment or when the dose is changed. Depression and other mental illnesses are the most important causes of suicidal thoughts and actions. Patients on antidepressants and their families or caregivers should watch for new or worsening depression symptoms, especially sudden changes in mood, behaviors, thoughts, or feelings. Report any change in these symptoms immediately to the doctor.

Please read [IMPORTANT SAFETY INFORMATION](#) on pages 9-11.



Continue to build on the progress you've made.

Actor portrayals



SELECT IMPORTANT SAFETY INFORMATION:

Do not take REXULTI if you are allergic to brexpiprazole or any of the ingredients in REXULTI.

REXULTI may cause serious side effects, including:

- Cerebrovascular problems, including stroke, in elderly people with dementia-related psychosis that can lead to death.
- Neuroleptic malignant syndrome (NMS) is a serious condition that can lead to death. Call your healthcare provider or go to the nearest hospital emergency room right away if you have some or all of the following signs and symptoms of NMS: high fever; changes in your pulse, blood pressure, heart rate, and breathing; stiff muscles; confusion; increased sweating

Please read [IMPORTANT SAFETY INFORMATION](#) on pages 9-11.



THE REXULTI SAVINGS CARD:

Easy to get.
Simple to use.



Text **SAVE** to 96747 to download the card directly to your phone.*
Show your card to your pharmacist and receive your savings plus refill reminders.

No activation required.†

Simply present your card to the pharmacist when you have your REXULTI prescription filled.
Or present it when having your generic ANTIDEPRESSANT filled.

Your pharmacist will keep your card on file for future use.

REXULTI

Two months of REXULTI as little as \$0 ✓

REXULTI Refills‡ as little as \$5 ✓

90-day Prescriptions as little as \$5 ✓

GENERIC ANTIDEPRESSANT (if prescribed)

Generic ANTIDEPRESSANT
as little as \$0 ✓

Up to **\$8,180 applied** to your annual deductible up to 12 uses†

Visit [REXULTISavings.com](https://www.rexulti.com/savings-card).

*Message and data rates apply. Message frequency varies. Text HELP for help and STOP to opt-out. Otsuka may use pharmacy information to provide refill reminders with your consent. Terms of Service: <https://rexulti.com/savings-card-terms-conditions>. Privacy Policy: <https://otsuka-us.com/oapi-and-opdc-privacy-policy>.

†Other terms and conditions apply, individual benefits may vary. See full terms and conditions at <https://www.rexulti.com/savings-card-terms-conditions>

‡This program is not health insurance. Further conditions apply. See full offer details on back of the card.

Program Eligibility

Offer is not transferable. Patients must have commercial drug insurance and a prescription consistent with FDA-approved product labeling. Patients are not eligible if they are covered in whole or in part by any state program or federal healthcare program, including, but not limited to, Medicare or Medicaid (including Medicaid managed care), Medigap, VA, DOD, or TRICARE. For eligible generic prescription, patient must be 18 years of age or older. Only valid in U.S. and Puerto Rico. Offer void where prohibited by law, taxed, or restricted. Other restrictions may apply. This program is not health insurance. Otsuka America Pharmaceutical, Inc. has the right to rescind, revoke, or amend this program at any time without notice. Your participation in this program confirms that this offer is consistent with your insurance coverage and that you will report the value received if required by your insurance provider. When you use this card, you are certifying that you understand and will comply with the program rules, terms, and conditions. Offer not valid for cash-paying patients OR where drug is not covered by the primary insurance.

Not eligible for the REXULTI Savings Card?

Learn about additional assistance. Call 1-855-242-7787, 8 AM–8 PM EST Monday through Friday, or visit otsukapatientssupport.com.

Still masking your **depression**?

You might be experiencing Partial Response.

Many people still experience depression symptoms, even on an antidepressant. Partial Response is when your antidepressant helps, but it's only part of what you need—and you're still left masking how you really feel inside.

Together with your doctor, there are things you can do to help with your depression symptoms, like adding REXULTI.

SELECT IMPORTANT SAFETY INFORMATION:

REXULTI® (brexpiprazole) may cause serious side effects, including: *(continued)*

- **Uncontrolled body movements (tardive dyskinesia).**
REXULTI may cause movements that you cannot control in your face, tongue, or other body parts. Tardive dyskinesia may not go away, even if you stop taking REXULTI. Tardive dyskinesia may also start after you stop taking REXULTI.
- **Problems with your metabolism such as:**
 - **high blood sugar (hyperglycemia) and diabetes.** Increases in blood sugar can happen in some people who take REXULTI. Extremely high blood sugar can lead to coma or death. Your healthcare provider should check your blood sugar before you start, or soon after you start REXULTI and then regularly during long term treatment with REXULTI.

Please read [IMPORTANT SAFETY INFORMATION](#) on pages 9-11.



Adding REXULTI® (brexpiprazole) has been proven to help reduce the symptoms of depression.



Actor portrayals

REXULTI works with your antidepressant to help build on any progress you've already made. So you may feel better—without feeling like you have to start over.

See clinical trial information on right for details.

SELECT IMPORTANT SAFETY INFORMATION:

Call your healthcare provider if you have any of these symptoms of high blood sugar during treatment with REXULTI:

- feel very thirsty
- feel very hungry
- feel sick to your stomach
- need to urinate more than usual
- feel weak or tired
- feel confused, or your breath smells fruity

Please read [IMPORTANT SAFETY INFORMATION](#) on pages 9-11.

REXULTI® (brexpiprazole) in clinical trials

REXULTI was studied in two 6-week clinical trials of 1,054 patients, aged 18-65, who had been diagnosed with major depressive disorder (MDD). The patients selected for the studies took an antidepressant for at least 2 months. Patients who then had ongoing symptoms of depression were randomly selected to receive 1 mg to 3 mg/day of REXULTI in addition to their antidepressant, while the rest continued taking their antidepressant alone.

At the end of the 6-week clinical trials, researchers discovered something: generally, adult patients whose doctors added REXULTI to their antidepressant showed more improvement than those who continued to take their antidepressant alone.

SELECT IMPORTANT SAFETY INFORMATION:

REXULTI may cause serious side effects, including: *(continued)*

- **Problems with your metabolism such as:** *(continued)*
 - **increased fat levels (cholesterol and triglycerides) in your blood.** Your healthcare provider should check the fat levels in your blood before you start, or soon after you start REXULTI, and then periodically during treatment with REXULTI.
 - **weight gain.** You and your healthcare provider should check your weight before you start and often during treatment with REXULTI.

How to take REXULTI[®] (brexpiprazole)

REXULTI is taken once daily, with or without food. It's important to ask your pharmacist or healthcare provider if you have any questions about taking REXULTI. Take REXULTI as your healthcare provider prescribed it:



- Do **not** change the dose or stop taking REXULTI yourself.
- Your healthcare provider may change your dose if needed. Do not change the dose or stop taking REXULTI without first talking to your healthcare provider.
- Take REXULTI 1 time each day with or without food.
- **Do not take REXULTI if you** are allergic to brexpiprazole or any of the ingredients in REXULTI. See end of Medication Guide for a complete list of ingredients in REXULTI.

If you take too much REXULTI, call your healthcare provider or the Poison Control Center at **1-800-222-1222** right away or go to the nearest hospital emergency room.

SELECT IMPORTANT SAFETY INFORMATION:

REXULTI may cause serious side effects, including: *(continued)*

- **Unusual and uncontrollable (compulsive) urges.** Some people taking REXULTI have had strong unusual urges, to gamble and gambling that cannot be controlled (compulsive gambling). Other compulsive urges include sexual urges, shopping, and eating or binge eating. If you or your family members notice that you are having new or unusual strong urges or behaviors, talk to your healthcare provider.
- **Low white blood cell count.** Your healthcare provider may do blood tests during the first few months of treatment with REXULTI.

Fill your prescription
for REXULTI
as prescribed by your
healthcare provider.

Watch a My Path Forward video to learn
about saving on your prescriptions.

[VIEW NOW](#)

My Path Forward[™]
VIDEO SERIES

Please read [IMPORTANT SAFETY INFORMATION](#) on pages 9-11.

Questions & Answers

Here are answers to a few of the questions that you may have about REXULTI.

Have questions that aren't answered here? Ask your healthcare provider at your next appointment.

Can I take other medications with REXULTI® (brexiprazole)?

Tell your healthcare provider about all the medicines you take or have recently taken, including prescription medicines, nonprescription medicines, vitamins, and herbal supplements. REXULTI and other medicines may affect each other, causing possible serious side effects. REXULTI may affect the way other medicines work, and other medicines may affect how REXULTI works.

Your healthcare provider can tell you if it's safe to take REXULTI with your other medicines. Don't start or stop any medicines without talking to your healthcare provider first. Only you and your healthcare provider can decide if REXULTI is right for you.

SELECT IMPORTANT SAFETY INFORMATION:

REXULTI may cause serious side effects, including: *(continued)*

- **Decreased blood pressure (orthostatic hypotension) and fainting.** You may feel dizzy, lightheaded or pass out (faint) when you rise too quickly from a sitting or lying position.

Please read [IMPORTANT SAFETY INFORMATION](#) on pages 9-11.



Questions & Answers

(continued)

When will REXULTI® (brexpiprazole) start working for me?

Not everyone responds to medication the same way, and your body may need time to adapt. That's why it's important to share how you are feeling with your healthcare provider, including any side effects. This way, he or she will better understand what effect your treatment may be having and make an adjustment if necessary. Always talk to your doctor before making any changes to your treatment.

What are the most common side effects?

The **most common side effects** associated with REXULTI plus an antidepressant were weight gain and restlessness or feeling like you need to move (akathisia), as reported during 6-week clinical studies of adults with major depressive disorder (MDD).

You and your healthcare provider should check your weight regularly. In 6-week clinical studies of adults with MDD, 7% of patients taking REXULTI plus an antidepressant reported an increase in body weight, compared with 2% of patients taking an antidepressant plus a placebo (nontherapeutic pill).

In the same studies, akathisia was reported by 9% of patients taking REXULTI plus an antidepressant and by 2% of patients taking an antidepressant plus a placebo.

SELECT IMPORTANT SAFETY INFORMATION:

REXULTI may cause serious side effects, including: (continued)

- **Falls.** REXULTI may make you sleepy or dizzy, may cause a decrease in your blood pressure when changing position (orthostatic hypotension), and can slow your thinking and motor skills which may lead to falls that can cause fractures or other injuries.
- **Seizures (convulsions).**

Please read [IMPORTANT SAFETY INFORMATION](#) on pages 9-11.



Actor portrayals

Questions & Answers

(continued)

Could REXULTI® (brexpiprazole) affect my blood sugar?

An increase in blood sugar (hyperglycemia) can happen in some people who take REXULTI. Extremely high blood sugar can lead to coma or death.

In clinical studies,* adult patients with shifts in fasting blood sugar from normal (<100 mg/dL) to high (≥ 126 mg/dL) and borderline (≥ 100 and <126 mg/dL) to high were similar in patients treated with REXULTI and patients treated with placebo.

Call your healthcare provider if you have any of these symptoms of high blood sugar while taking REXULTI: You feel very thirsty; feel very hungry; feel sick to your stomach; need to urinate more than usual; feel weak or tired; feel confused, or your breath smells fruity.

Will REXULTI affect my cholesterol?

An increase in the fat levels (cholesterol and triglycerides) in your blood may occur while taking REXULTI. Before starting on REXULTI, discuss with your healthcare provider if you have high levels of cholesterol, triglycerides, or LDL cholesterol, or low levels of HDL cholesterol.

Your healthcare provider should check the fat levels in your blood before you start, or soon after you start REXULTI, and then periodically during treatment with REXULTI.

During clinical trials,* changes in fasting total cholesterol, LDL cholesterol, and HDL cholesterol were similar between adult patients taking REXULTI and patients taking a placebo.

Will REXULTI make me feel sleepy or drowsy?

In clinical trials,* 4% of adult patients who had REXULTI added to an antidepressant experienced drowsiness, compared with 1% of patients taking an antidepressant plus a placebo.

Until you know how REXULTI affects you, do not drive, operate heavy machinery, or do dangerous activities.

Will REXULTI make me feel restless?

About 3% of adult patients taking REXULTI plus an antidepressant reported feelings of restlessness during clinical trials,* compared with 0% of patients taking placebo.

Does REXULTI affect sexual function?

Overall, there was no difference seen in sexual function of patients as a result of taking REXULTI with an antidepressant compared with those taking an antidepressant alone.

During clinical trials,* adult patients taking REXULTI plus an antidepressant reported their overall sexual function was similar to that of patients taking an antidepressant alone, as measured by the Massachusetts General Hospital Sexual Functioning Questionnaire.

*Two 6-week clinical trials. For more information about the clinical trials, please see page 4.

SELECT IMPORTANT SAFETY INFORMATION:

REXULTI may cause serious side effects, including: (continued)

- **Problems controlling your body temperature so that you feel too warm.** Do not become too hot or dehydrated during treatment with REXULTI. Do not exercise too much. In hot weather, stay inside in a cool place if possible. Stay out of the sun. Do not wear too much clothing or heavy clothing. Drink plenty of water.

Please read [IMPORTANT SAFETY INFORMATION](#) on pages 9-11.

INDICATION and IMPORTANT SAFETY INFORMATION for REXULTI® (brexpiprazole)

INDICATION: REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

SELECT IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis without agitation that may happen with dementia due to Alzheimer's disease.
- Increased risk of suicidal thoughts and actions. REXULTI and antidepressant medicines may increase suicidal thoughts and actions in some people 24 years of age and younger, especially within the first few months of treatment or when the dose is changed. Depression and other mental illnesses are the most important causes of suicidal thoughts and actions. Patients on antidepressants and their families or caregivers should watch for new or worsening depression symptoms, especially sudden changes in mood, behaviors, thoughts, or feelings. Report any change in these symptoms immediately to the doctor.

Do not take REXULTI if you are allergic to brexpiprazole or any of the ingredients in REXULTI.

(Continued on next page)

Please read **FULL PRESCRIBING INFORMATION**, including **BOXED WARNING**, and **MEDICATION GUIDE**.



IMPORTANT SAFETY INFORMATION

(continued)

REXULTI® (brexpiprazole) may cause serious side effects, including:

- **Cerebrovascular problems, including stroke, in elderly people with dementia-related psychosis that can lead to death.**
- **Neuroleptic malignant syndrome (NMS) is a serious condition that can lead to death.** Call your healthcare provider or go to the nearest hospital emergency room right away if you have some or all of the following signs and symptoms of NMS: high fever; changes in your pulse, blood pressure, heart rate, and breathing; stiff muscles; confusion; increased sweating

- **Uncontrolled body movements (tardive dyskinesia).**

REXULTI may cause movements that you cannot control in your face, tongue, or other body parts. Tardive dyskinesia may not go away, even if you stop taking REXULTI. Tardive dyskinesia may also start after you stop taking REXULTI.

- **Problems with your metabolism such as:**

- **high blood sugar (hyperglycemia) and diabetes.**

Increases in blood sugar can happen in some people who take REXULTI. Extremely high blood sugar can lead to coma or death. Your healthcare provider should check your blood sugar before you start, or soon after you start REXULTI and then regularly during long term treatment with REXULTI.

Call your healthcare provider if you have any of these symptoms of high blood sugar during treatment with REXULTI:

- feel very thirsty
 - feel very hungry
 - feel sick to your stomach
 - need to urinate more than usual
 - feel weak or tired
 - feel confused, or your breath smells fruity

- **increased fat levels (cholesterol and triglycerides) in your blood.** Your healthcare provider should check the fat levels in your blood before you start, or soon after you start REXULTI, and then periodically during treatment with REXULTI.

- **weight gain.** You and your healthcare provider should check your weight before you start and often during treatment with REXULTI.

- **Unusual and uncontrollable (compulsive) urges.** Some people taking REXULTI have had strong unusual urges, to gamble and gambling that cannot be controlled (compulsive gambling). Other compulsive urges include sexual urges, shopping, and eating or binge eating. If you or your family members notice that you are having new or unusual strong urges or behaviors, talk to your healthcare provider.

- **Low white blood cell count.** Your healthcare provider may do blood tests during the first few months of treatment with REXULTI.

- **Decreased blood pressure (orthostatic hypotension) and fainting.** You may feel dizzy, lightheaded or pass out (faint) when you rise too quickly from a sitting or lying position.

- **Falls.** REXULTI may make you sleepy or dizzy, may cause a decrease in your blood pressure when changing position (orthostatic hypotension), and can slow your thinking and motor skills which may lead to falls that can cause fractures or other injuries.

- **Seizures (convulsions).**

- **Problems controlling your body temperature so that you feel too warm.** Do not become too hot or dehydrated during treatment with REXULTI. Do not exercise too much. In hot weather, stay inside in a cool place if possible. Stay out of the sun. Do not wear too much clothing or heavy clothing. Drink plenty of water.

- **Difficulty swallowing** that can cause food or liquid to get into your lungs.

- **Sleepiness, drowsiness, feeling tired, difficulty thinking and doing normal activities.** Do not drive a car, operate machinery, or do other dangerous activities until you know how REXULTI affects you. REXULTI may make you feel drowsy.

(Continued on next page)

Please read **FULL PRESCRIBING INFORMATION**, including **BOXED WARNING**, and **MEDICATION GUIDE**.

IMPORTANT SAFETY INFORMATION

(continued)

Before taking REXULTI® (brexpiprazole), tell your healthcare provider about all of your medical conditions, including if you:

- have or have had heart problems or a stroke
- have or have had low or high blood pressure
- have or have had diabetes or high blood sugar or a family history of diabetes or high blood sugar. Your healthcare provider should check your blood sugar before you start REXULTI and during treatment with REXULTI.
- have or have had high levels of total cholesterol, LDL cholesterol, or triglycerides, or low levels of HDL cholesterol
- have or have had seizures (convulsions)
- have or have had kidney or liver problems
- have or have had a low white blood cell count
- are pregnant or plan to become pregnant. REXULTI may harm your unborn baby. Taking REXULTI during your third trimester of pregnancy may cause your baby to have abnormal muscle movements or withdrawal symptoms after birth. Talk to your healthcare provider about the risk to your unborn baby if you take REXULTI during pregnancy.
 - Tell your healthcare provider if you become pregnant or think you are pregnant during treatment with REXULTI.
 - There is a pregnancy exposure registry for women who are exposed to REXULTI during pregnancy. If you become pregnant during treatment with REXULTI, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics. You can register by calling 1-866-961-2388 or visit <http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/>.
- are breastfeeding or plan to breastfeed. It is not known if REXULTI passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with REXULTI.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. REXULTI and other medicines may affect each other causing possible serious side effects. REXULTI may affect the way other medicines work, and other medicines may affect how REXULTI works. Your healthcare provider can tell you if it is safe to take REXULTI with your other medicines. Do not start or stop any medicines during treatment with REXULTI without first talking to your healthcare provider.

The most common side effects of REXULTI include weight gain, sleepiness, dizziness, common cold symptoms, and restlessness or feeling like you need to move (akathisia).

These are not all the possible side effects of REXULTI. For more information, ask your healthcare provider or pharmacist.

You are encouraged to report side effects of REXULTI (brexpiprazole). Please contact Otsuka America Pharmaceutical, Inc. at 1-800-438-9927 or FDA at 1-800-FDA-1088 (www.fda.gov/medwatch).

Please read **FULL PRESCRIBING INFORMATION**, including **BOXED WARNING**, and **MEDICATION GUIDE**.

Resources for every step of your journey

Please read [FULL PRESCRIBING INFORMATION](#), including [BOXED WARNING](#), and [MEDICATION GUIDE](#).



Watch the My Path Forward videos for help on:

- Understanding your symptoms
- Talking to your doctor
- Saving on prescriptions

[CLICK HERE TO WATCH](#) ➤



Use this online guide to help with your overall wellness.

My Wellness Guide was created especially for you. Full of handy tips and tools, simply download the book and refer to it as needed.

[CLICK HERE TO DOWNLOAD](#) ➤



Get your complimentary journal.

Track your progress with tips and tools.

[CLICK HERE TO GET YOUR JOURNAL](#) ➤

For more information visit [REXULTI.com](https://www.REXULTI.com).