The REXULTI Savings Card Frequently Asked Questions







What is different with the new REXULTI Savings Card?

The REXULTI Savings Card

now covers a generic antidepressant too.

For new and existing REXULTI and generic antidepressant patients:

- \$0 for the first 2 months
- Maximum applied for first two months is \$1,300/month (\$2,600 total)
- Pay as little as \$15 per refill
- Maximum applied per refill is \$200/month
- Pay as little as \$15 for 90-day prescriptions
- 1 card/patient in calendar year
- \$0 per month for generic antidepressant with maximum applied of \$200/month

Does the REXULTI Savings Card need to be activated?

No, the REXULTI Savings Card is ready to use and requires no activation.

When does the REXULTI Savings Card expire?

There is no expiration date for the REXULTI Savings Card and no current plans to discontinue this offer. Otsuka America Pharmaceutical, Inc. has the right to rescind, revoke, or amend this program at any time without notice.

Are there any restrictions by insurance and/or pharmacy?

Patients may be eligible for the Savings Card Offer if they have commercial prescription insurance and are 18-64 years of age. Full eligibility will be determined by their pharmacist. For reference, REXULTI is covered on 97% of commercial insurance plans nationwide. Most pharmacies accept savings cards. In the rare instance where a pharmacy does not allow for copay card use, a patient can request reimbursement by downloading a form from rexulti.com/mailorder.

If a patient's insurance changes, can they still use their REXULTI Savings Card?

As long as the new insurance covers REXULTI and/or the generic antidepressant and is not Medicare or Medicaid (including managed Medicaid), Medigap, VA, DOD, or TRICARE, or another federally or state funded program that covers their prescription in whole or in part; and this offer is consistent with their insurance, they should be able to use the REXULTI Savings Card.

INDICATION:

REXULTI is a prescription medicine used:

• with other antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

SELECT IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis.
- Increased risk of suicidal thoughts and actions. Antidepressant medicines may increase suicidal thoughts and actions in some children, adolescents, and young adults especially within the first few months of treatment or when the dose is changed. Depression and other mental illnesses are the most important causes of suicidal thoughts and actions.
 Patients on antidepressants and their families or caregivers should watch for new or worsening depression symptoms, especially sudden changes in mood, behaviors, thoughts, or feelings. Report any change in these symptoms immediately to the doctor.

What if the patient has a deductible vs a copay?

If REXULTI® (brexpiprazole) and/or the generic antidepressant is covered by the patient's insurance, both the patient's out-of-pocket and co-pay card benefit will count towards their deductible.

For the first 2 uses, patients pay \$0 for REXULTI with a maximum benefit of \$1,300 per use—\$2,600 that may be applied to an annual deductible.

For uses 3+, patients may pay as little as \$15 for REXULTI with a maximum benefit of \$200 per use. That is an additional \$200 per month that may be applied to an annual deductible.

For the generic antidepressant, patients may pay \$0 per use with a maximum benefit of \$100.

Up to \$5,800 may be applied to an annual deductible with 12 uses. If a patient's deductible has already been met, a patient's expected out-of-pocket cost still operates as outlined above.

Do patients need to acquire both medications from the same pharmacy?

No, but patients will need to present the same card for use at both pharmacies and ask the pharmacist to retain the card information on file. Once the card information is on file for both prescriptions, they should not need to bring it each time they fill a prescription.

What if they are on a branded antidepressant instead of a generic?

The REXULTI Savings Card does not cover branded antidepressants.

What if they are on more than one generic antidepressant?

The REXULTI Savings Card will cover one generic antidepressant in a 30-day period.

What generic antidepressants will the REXULTI Savings Card cover?

The REXULTI Savings Card covers currently available generic antidepressants. Please see **REXULTI.com** for a list of covered generic antidepressants.

Can patients use the card for the antidepressant alone?

The benefits for REXULTI and the generic antidepressant are separate. The generic antidepressant can be written within 3 months before and after treatment with REXULTI.

• Those patients can still receive their separate benefits for both prescriptions within that proximate timeframe.

Can patients still use the card for their generic antidepressant if they stop taking REXULTI?

The benefits for REXULTI and the generic antidepressant are separate. The generic antidepressant can be written within 3 months before and after treatment with REXULTI.

• Those patients can still receive their separate benefits for both prescriptions within that proximate timeframe.

Does the REXULTI Savings Card work for 90-day prescriptions?

Yes, patients should confirm with their insurance providers. If allowed, patients pay as little as \$15 per 90-day prescriptions for REXULTI and \$0 for the generic antidepressant.

What if a patient is already taking REXULTI?

An eligible, existing REXULTI patient can still use the REXULTI Savings Card for REXULTI refills and generic antidepressant prescriptions. Patients should present the card to the pharmacist when picking up their prescription(s) and ask the pharmacist to keep the card information on file. The \$0 for 2 months offer is only valid for one use in a calendar year. If an existing REXULTI patient used a previous REXULTI Savings Card in the same calendar year, they will still be eligible to pay as little as \$15 per refill for REXULTI and \$0 for a generic antidepressant.

Dose the REXULTI Savings Card apply to all doses?

Yes, the REXULTI Savings Card applies to all doses of REXULTI.

SELECT IMPORTANT SAFETY INFORMATION:

Do not take REXULTI if you are allergic to brexpiprazole or any of the ingredients in REXULTI.

INDICATION and IMPORTANT SAFETY INFORMATION for REXULTI® (brexpiprazole) INDICATION:

REXULTI is a prescription medicine used:

• with other antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis.
- Increased risk of suicidal thoughts and actions. Antidepressant medicines may increase suicidal thoughts and actions in some children, adolescents, and young adults especially within the first few months of treatment or when the dose is changed. Depression and other mental illnesses are the most important causes of suicidal thoughts and actions.
 Patients on antidepressants and their families or caregivers should watch for new or worsening depression symptoms, especially sudden changes in mood, behaviors, thoughts, or feelings. Report any change in these symptoms immediately to the doctor.

Do not take REXULTI if you are allergic to brexpiprazole or any of the ingredients in REXULTI.

REXULTI may cause serious side effects, including:

- Stroke (cerebrovascular problems) in elderly people with dementia-related psychosis that can lead to death.
- Neuroleptic malignant syndrome (NMS) is a serious condition that can lead to death. Call your healthcare provider or go to the nearest hospital emergency room right away if you have some or all of the following signs and symptoms of NMS: high fever; changes in your breathing, heart rate, and blood pressure; stiff muscles; confusion; increased sweating.
- **Uncontrolled body movements (tardive dyskinesia).** REXULTI may cause movements that you cannot control in your face, tongue, or other body parts. Tardive dyskinesia may not go away, even if you stop taking REXULTI. Tardive dyskinesia may also start after you stop taking REXULTI.
- Problems with your metabolism such as:
 - o high blood sugar (hyperglycemia) and diabetes: Increases in blood sugar can happen in some people who take REXULTI. Extremely high blood sugar can lead to coma or death. Your healthcare provider should check your blood sugar before you start, or soon after you start REXULTI and then regularly during long term treatment with REXULTI.

Call your healthcare provider if you have any of these symptoms of high blood sugar during treatment with REXULTI:

- feel very thirsty
- feel very hungry
- feel sick to your stomach
- need to urinate more than usual
- feel weak or tired
- feel confused, or your breath smells fruity
- o increased fat levels (cholesterol and triglycerides) in your blood. Your healthcare provider should check the fat levels in your blood before you start, or soon after you start REXULTI, and then periodically during treatment with REXULTI.
- oweight gain. You and your healthcare provider should check your weight before you start and often during treatment with REXULTI.
- Unusual and uncontrollable (compulsive) urges. Some people taking REXULTI have had strong unusual urges, to gamble and gambling that cannot be controlled (compulsive gambling). Other compulsive urges include sexual urges, shopping, and eating or binge eating. If you or your family members notice that you are having unusual strong urges, talk to your healthcare provider.
- Low white blood cell count. Your healthcare provider may do blood tests during the first few months of treatment with REXULTI.
- Decreased blood pressure (orthostatic hypotension). You may feel lightheaded or faint when you rise too quickly from a sitting or lying position.
- Falls. REXULTI may make you sleepy or dizzy, may cause a decrease in your blood pressure when changing position (orthostatic hypotension), and can slow your thinking and motor skills which may lead to falls that can cause fractures or other injuries.
- Seizures (convulsions).
- Problems controlling your body temperature so that you feel too warm. Do not become too hot or dehydrated during treatment with REXULTI. Do not exercise too much. In hot weather, stay inside in a cool place if possible. Stay out of the sun. Do not wear too much clothing or heavy clothing. Drink plenty of water.
- Difficulty swallowing that can cause food or liquid to get into your lungs.
- Sleepiness, drowsiness, feeling tired, difficulty thinking and doing normal activities. Do NOT drive a car, operate machinery, or do other dangerous activities until you know how REXULTI affects you. REXULTI may make you feel drowsy.

(Continued on next page)

IMPORTANT SAFETY INFORMATION for REXULTI® (brexpiprazole) (continued)

Before taking REXULTI, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had heart problems or a stroke
- have or have had low or high blood pressure
- have or have had diabetes or high blood sugar or a family history of diabetes or high blood sugar. Your healthcare provider should check your blood sugar before you start REXULTI and during treatment with REXULTI.
- have or have had high levels of total cholesterol, LDL cholesterol, or triglycerides, or low levels of HDL cholesterol
- have or have had seizures (convulsions)
- have or have had kidney or liver problems
- have or have had a low white blood cell count
- are pregnant or plan to become pregnant. REXULTI may harm your unborn baby. Taking REXULTI during your third trimester of pregnancy may cause your baby to have abnormal muscle movements or withdrawal symptoms after birth.
 - Tell your healthcare provider if you become pregnant or think you are pregnant during treatment with REXULTI.
 - o If you become pregnant during treatment with REXULTI, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics. You can register by calling 1-866-961-2388 or visit http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/.
- are breastfeeding or plan to breastfeed. It is not known if REXULTI passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with REXULTI.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. REXULTI and other medicines may affect each other causing possible serious side effects. REXULTI may affect the way other medicines work, and other medicines may affect how REXULTI works. Your healthcare provider can tell you if it is safe to take REXULTI with your other medicines. Do not start or stop any medicines during treatment with REXULTI without first talking to your healthcare provider.

The most common side effects of REXULTI include weight gain and restlessness or feeling like you need to move (akathisia).

These are not all the possible side effects of REXULTI. For more information, ask your healthcare provider or pharmacist.

You are encouraged to report side effects of REXULTI (brexpiprazole). Please contact Otsuka America Pharmaceutical, Inc. at 1-800-438-9927 or FDA at 1-800-FDA-1088 (<u>www.fda.gov/medwatch</u>).

Please read FULL PRESCRIBING INFORMATION, including BOXED WARNING, and MEDICATION GUIDE.



