

Here's Your **REXULTI** Savings Card:

Using the card is **SIMPLE.**

No activation required.

Simply present your card to the pharmacist when you have your REXULTI or your generic antidepressant filled (if prescribed).

Your pharmacist will keep card on file for future use.



Offer not valid for cash-paying patients OR where drug is not covered by primary insurance.

REXULTI

Two months of* REXULTI as little as

REXULTI refills† as little as

90-day prescriptions as little as

GENERIC ANTIDEPRESSANT (if prescribed)

Generic ANTIDEPRESSANT as little as

\$0



Up to \$8,180 APPLIED to your annual deductible up to 12 uses*

INDICATIONS AND IMPORTANT SAFETY INFORMATION FOR REXULTI® (brexpiprazole) INDICATIONS:

REXULTI is a prescription medicine used:

- along with antidepressant medicines to treat major depressive disorder (MDD) in adults
- to treat schizophrenia in adults and children ages 13 years and older
- to treat agitation that may happen with dementia due to Alzheimer's disease

REXULTI should not be used as an "as needed" treatment for agitation that may happen with dementia due to Alzheimer's disease. It is not known if REXULTI is safe and effective in children with MDD.

It is not known if REXULTI is safe and effective in children under 13 years of age with schizophrenia.

SELECT IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in
 elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved
 for the treatment of people with dementia-related psychosis without agitation that may happen with dementia due to Alzheimer's
 disease.
- Increased risk of suicidal thoughts and actions. REXULTI and antidepressant medicines may increase suicidal thoughts and actions in some people 24 years of age and younger, especially within the first few months of treatment or when the dose is changed. Depression and other mental illnesses are the most important causes of suicidal thoughts and actions. Patients on antidepressants and their families or caregivers should watch for new or worsening depression symptoms, especially sudden changes in mood, behaviors, thoughts, or feelings. Report any change in these symptoms immediately to the doctor.

Do not take REXULTI if you are allergic to brexpiprazole or any of the ingredients in REXULTI.

^{*} Offer not valid for cash-paying patients OR where drug is not covered by primary insurance.

[†]This program is not health insurance. Further conditions apply. See full offer details on page 3.

IMPORTANT SAFETY INFORMATION FOR REXULTI® (brexpiprazole) (continued)

REXULTI may cause serious side effects, including:

- Cerebrovascular problems, including stroke, in elderly people with dementia-related psychosis that can lead to death.
- Neuroleptic malignant syndrome (NMS) is a serious condition that can lead to death. Call your healthcare provider or go to the nearest hospital emergency room right away if you have some or all of the following signs and symptoms of NMS: high fever; changes in your pulse, blood pressure, heart rate, and breathing; stiff muscles; confusion; increased sweating.
- Uncontrolled body movements (tardive dyskinesia). REXULTI may cause movements that you cannot control in your face, tongue, or other body parts. Tardive dyskinesia may not go away, even if you stop taking REXULTI. Tardive dyskinesia may also start after you stop taking REXULTI.
- Problems with your metabolism such as:
 - o high blood sugar (hyperglycemia) and diabetes. Increases in blood sugar can happen in some people who take REXULTI. Extremely high blood sugar can lead to coma or death. Your healthcare provider should check your blood sugar before you start, or soon after you start REXULTI and then regularly during long term treatment with REXULTI.

Call your healthcare provider if you have any of these symptoms of high blood sugar during treatment with REXULTI:

- feel very thirsty
- feel very hungry
- feel sick to your stomach
- need to urinate more than usual
- feel weak or tired
- feel confused, or your breath smells fruity
- o increased fat levels (cholesterol and triglycerides) in your blood. Your healthcare provider should check the fat levels in your blood before you start, or soon after you start REXULTI, and then periodically during treatment with REXULTI.
- weight gain. You and your healthcare provider should check your weight before you start and often during treatment with REXULTI.
- Unusual and uncontrollable (compulsive) urges. Some people
 taking REXULTI have had strong unusual urges, to gamble and
 gambling that cannot be controlled (compulsive gambling). Other
 compulsive urges include sexual urges, shopping, and eating or
 binge eating. If you or your family members notice that you are
 having new or unusual strong urges or behaviors, talk to your
 healthcare provider.
- Low white blood cell count. Your healthcare provider may do blood tests during the first few months of treatment with REXULTI.
- Decreased blood pressure (orthostatic hypotension) and fainting. You may feel dizzy, lightheaded or pass out (faint) when you rise too quickly from a sitting or lying position.
- Falls. REXULTI may make you sleepy or dizzy, may cause a decrease in your blood pressure when changing position (orthostatic hypotension), and can slow your thinking and motor skills which may lead to falls that can cause fractures or other injuries.
- Seizures (convulsions).
- Problems controlling your body temperature so that you feel too warm. Do not become too hot or dehydrated during treatment

- with REXULTI. Do not exercise too much. In hot weather, stay inside in a cool place if possible. Stay out of the sun. Do not wear too much clothing or heavy clothing. Drink plenty of water.
- **Difficulty swallowing** that can cause food or liquid to get into your lungs.
- Sleepiness, drowsiness, feeling tired, difficulty thinking and doing normal activities. Do not drive a car, operate machinery, or do other dangerous activities until you know how REXULTI affects you.
 REXULTI may make you feel drowsy.

Before taking REXULTI, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had heart problems or a stroke
- have or have had low or high blood pressure
- have or have had diabetes or high blood sugar or a family history
 of diabetes or high blood sugar. Your healthcare provider should
 check your blood sugar before you start REXULTI and during
 treatment with REXULTI.
- have or have had high levels of total cholesterol, LDL cholesterol, or triglycerides, or low levels of HDL cholesterol
- have or have had seizures (convulsions)
- have or have had kidney or liver problems
- have or have had a low white blood cell count
- are pregnant or plan to become pregnant. REXULTI may harm your unborn baby. Taking REXULTI during your third trimester of pregnancy may cause your baby to have abnormal muscle movements or withdrawal symptoms after birth. Talk to your healthcare provider about the risk to your unborn baby if you take REXULTI during pregnancy.
 - Tell your healthcare provider if you become pregnant or think you are pregnant during treatment with REXULTI.
 - There is a pregnancy exposure registry for women who are exposed to REXULTI during pregnancy. If you become pregnant during treatment with REXULTI, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics. You can register by calling 1-866-961-2388 or visit http:// womensmentalhealth.org/clinical-and-researchprograms/pregnancyregistry/.
- are breastfeeding or plan to breastfeed. It is not known if REXULTI passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with REXULTI.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. REXULTI and other medicines may affect each other causing possible serious side effects. REXULTI may affect the way other medicines work, and other medicines may affect how REXULTI works. Your healthcare provider can tell you if it is safe to take REXULTI with your other medicines. Do not start or stop any medicines during treatment with REXULTI without first talking to your healthcare provider.

The most common side effects of REXULTI include weight gain, sleepiness, dizziness, common cold symptoms, and restlessness or feeling like you need to move (akathisia).

These are not all the possible side effects of REXULTI. For more information, ask your healthcare provider or pharmacist.

You are encouraged to report side effects of REXULTI (brexpiprazole). Please contact Otsuka America Pharmaceutical, Inc. at

1-800-438-9927 or FDA at 1-800-FDA-1088

(www.fda.gov/medwatch).

For help using this card, call 1-844-415-0674.

Terms, Conditions, and Eligibility Criteria: This program is offered by Otsuka America Pharmaceutical, Inc. and applies only to valid REXULTI® (brexpiprazole) and/or eligible generic prescriptions. Eligible insured patients will pay as little as \$0 for first two uses 8-33 and 34-60-day supply and as little as \$5 for 90-day supply. On subsequent uses 3 and beyond, patients will pay as little as \$5 and receive a maximum savings up to \$1,200. For eligible generic prescription with 30-day supply patient will pay as little as \$0 and receive savings up to \$100 per 30-day supply. **Up to \$8,180 applied to the annual deductible with up to 12 uses.** Limit 1 card per patient.

Patient Instructions: In order to redeem this offer you must have a valid prescription for REXULTI and/or an eligible generic prescription. Follow the dosage instructions given by the doctor. This offer may not be redeemed for cash. By using this offer, you are certifying that you meet the eligibility criteria and will comply with the terms and conditions described in the Restrictions section below.

Pharmacist: When you use this card, you are certifying that you have not submitted and will not submit a claim for reimbursement under any federal, state, or other governmental programs for this prescription. As a condition of payment, you certify that you are in compliance with all program rules, terms, and conditions, as well as with any obligations to provide notice of your participation in this program to third-party payers as required by law, contract, or otherwise.

Submit the claim to the primary Third-Party Payer first, then submit the balance due to PDMI as a Secondary Payer COB [coordination of benefits] with patient responsibility amount and a valid Other Coverage Code (eg. 8).

For days' supply of 7 days or less, the patient is responsible for a \$0 copay. Reimbursement will be received from PDMI. For any questions regarding PDMI online processing, please call the Help Desk at 1-833-742-0793.

Restrictions: Offer is not transferable. Patients must have commercial drug insurance and a prescription consistent with FDA-approved product labeling. Patients are not eligible if they are covered in whole or in part by any state program or federal healthcare program, including, but not limited to, Medicare or Medicaid (including Medicaid managed care), Medigap, VA, DOD, or TRICARE. For eligible generic prescription, patient must be 18 years of age or older. Only valid in U.S. and Puerto Rico. Offer void where prohibited by law, taxed, or restricted. Other restrictions may apply. This program is not health insurance. Otsuka America Pharmaceutical, Inc. has the right to rescind, revoke, or amend this program at any time without notice. Your participation in this program confirms that this offer is consistent with your insurance coverage and that you will report the value received if required by your insurance provider. When you use this card, you are certifying that you understand and will comply with the program rules, terms, and conditions. Offer not valid for cash-paying patients OR where drug is not covered by the

Please read FULL PRESCRIBING INFORMATION, including BOXED WARNING, and MEDICATION GUIDE.



