REXULTI® (brexpiprazole) Savings Program Patient Reimbursement

Reimbursement through the REXULTI Savings Program for out-of-pocket co-pay cost directly incurred by the patient/caregiver for REXULTI or a Generic Antidepressant will be available to eligible applicants who meet the eligibility requirements listed below (see full terms and conditions).

To receive reimbursement, please complete the accompanying Reimbursement Form. Submit the form and all required documentation to:

Rx Savings Program PO Box 2355 Morristown, NJ 07962 or

Fax: 1-908-809-6230

REXULTI Savings Program Terms and Conditions

Program Eligibility:

This program is offered by Otsuka America Pharmaceutical, Inc. and applies only to valid REXULTI® (brexpiprazole) and/or eligible generic prescriptions. Eligible insured patients will pay \$0 for first two uses 8-33 and 34-60-day supply and as little as \$15 for 90-day supply. On subsequent uses 3 and beyond, patients will pay as little as \$15 and receive a maximum saving up to \$600. For a valid eligible generic prescription with 30-day supply patient will pay as little as \$0 and receive savings up to \$100 per use. Limit 1 card per patient.

Patient Instructions:

In order to redeem this offer you must have a valid prescription for REXULTI or a Generic Antidepressant. Follow the dosage instructions given by the doctor. This offer may not be redeemed for cash. By using this offer, you are certifying that you meet the eligibility criteria and will comply with the terms and conditions described in the Restrictions section below. Patients with questions about the REXULTI savings offers should call 1-844-415-0674.

Restrictions:

Offers are not transferable. Patients must have commercial drug insurance and a prescription consistent with FDA-approved product labeling. Patients are not eligible if they are covered in whole or in part by any state program or federal healthcare program, including, but not limited to, Medicare or Medicaid (including Medicaid managed care), Medigap, VA, DOD, or TRICARE. For eligible generic prescription, patient must be 18 years of age or older. Only valid in U.S. and Puerto Rico. Offers void where prohibited by law, taxed, or restricted. Other restrictions may apply. This program is not health insurance. Otsuka America Pharmaceutical, Inc. has the right to rescind, revoke, or amend this program at any time without notice. Your participation in this program confirms that the offer is consistent with your insurance coverage and that you will report the value received if required by your insurance provider. When you use this card, you are certifying that you understand and will comply with the program rules, terms, and conditions. Offers not valid for cash-paying patients OR where drug is not covered by the primary insurance.

INDICATION:

REXULTI is a prescription medicine used:

• with other antidepressant to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

SELECT IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death
 in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not
 approved for the treatment of people with dementia-related psychosis.
- Increased risk of suicidal thoughts and actions. Antidepressant medicines may increase suicidal thoughts and actions in some children, adolescents, and young adults especially within the first few months of treatment or when the dose is changed. Depression and other mental illnesses are the most important causes of suicidal thoughts and actions. Patients on antidepressants and their families or caregivers should watch for new or worsening depression symptoms, especially sudden changes in mood, behaviors, thoughts, or feelings. Report any change in these symptoms immediately to the doctor.

Please print all information and mail or fax all required materials to:

Rx Savings Program PO Box 2355 Morristown, NJ 07962 or

Fax: 1-908-809-6230

This form is to be used for reimbursement of a patient's co-pay out-of-pocket expenses directly and actually incurred for a prescribed dose of REXULTI® (brexpiprazole) or a Generic Antidepressant under the REXULTI Savings Program sponsored by Otsuka America Pharmaceutical, Inc. The REXULTI Savings Program reimburses eligible patients for their REXULTI or Generic Antidepressant product co-pay cost. Costs incurred for general office visits will not be reimbursed. Payment of the reimbursement is subject to verification.

	PATIENT INFOR	RMATION - pleas	se print	
First Name	Middle		Last Name	
Address 1		Address 2		
City		State	Zip	
Phone		Email		
DOB	Gende	er	Age	
	PRESCRIBER INFO	<u>ORMATION – pl</u>	ease print	
First Name	Middle		Last Name	
Address 1		Address 2		
ty		State	Zip	
Phone				
 Copy of front and Copy of REXULT days supply) Dated receipt (prexpenses highlig Patient signature 	e to receive reimbursement. For d back of your REXULTI Savings of or Generic Antidepressant preson of purchase or invoice) with thed	ms will take one to Card. scription label (inc the amount of co	ludes name and address of pharm	acy, dosing, and
accurate to the best of my kno		pocket expenses	requested for reimbursement were	
			/site/privacy-policy to learn more.	

SELECT IMPORTANT SAFETY INFORMATION:

Do not take REXULTI if you are allergic to brexpiprazole or any of the ingredients in REXULTI.

Otsuka's privacy policy can be found at https://www.otsuka-us.com/oapi-and-opdc-privacy-policy.

INDICATION and IMPORTANT SAFETY INFORMATION for REXULTI® (brexpiprazole) INDICATION:

REXULTI is a prescription medicine used:

• with other antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of
 death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia).
 REXULTI is not approved for the treatment of people with dementia-related psychosis.
- Increased risk of suicidal thoughts and actions. Antidepressant medicines may increase suicidal thoughts and actions
 in some children, adolescents, and young adults especially within the first few months of treatment or when the dose is
 changed. Depression and other mental illnesses are the most important causes of suicidal thoughts and actions.
 Patients on antidepressants and their families or caregivers should watch for new or worsening depression symptoms,
 especially sudden changes in mood, behaviors, thoughts, or feelings. Report any change in these symptoms
 immediately to the doctor.

Do not take REXULTI if you are allergic to brexpiprazole or any of the ingredients in REXULTI.

REXULTI may cause serious side effects, including:

- Stroke (cerebrovascular problems) in elderly people with dementia-related psychosis that can lead to death.
- Neuroleptic malignant syndrome (NMS) is a serious condition that can lead to death. Call your healthcare provider or go to the nearest hospital emergency room right away if you have some or all of the following signs and symptoms of NMS: high fever; changes in your breathing, heart rate, and blood pressure; stiff muscles; confusion; increased sweating.
- Uncontrolled body movements (tardive dyskinesia). REXULTI may cause movements that you cannot control in your face, tongue, or other body parts. Tardive dyskinesia may not go away, even if you stop taking REXULTI. Tardive dyskinesia may also start after you stop taking REXULTI.
- Problems with your metabolism such as:
 - high blood sugar (hyperglycemia) and diabetes. Increases in blood sugar can happen in some people who take REXULTI. Extremely high blood sugar can lead to coma or death. Your healthcare provider should check your blood sugar before you start, or soon after you start REXULTI and then regularly during long term treatment with REXULTI.

Call your healthcare provider if you have any of these symptoms of high blood sugar during treatment with REXULTI:

- feel very thirsty
- feel very hungry
- feel sick to your stomach
- need to urinate more than usual
- feel weak or tired
- feel confused, or your breath smells fruity
- o increased fat levels (cholesterol and triglycerides) in your blood. Your healthcare provider should check the fat levels in your blood before you start, or soon after you start REXULTI, and then periodically during treatment with REXULTI.
- o weight gain. You and your healthcare provider should check your weight before you start and often during treatment with REXULTI.
- Unusual and uncontrollable (compulsive) urges. Some people taking REXULTI have had strong unusual urges, to gamble and gambling that cannot be controlled (compulsive gambling). Other compulsive urges include sexual urges, shopping, and eating or binge eating. If you or your family members notice that you are having unusual strong urges, talk to your healthcare provider.
- Low white blood cell count. Your healthcare provider may do blood tests during the first few months of treatment with REXULTI.
- Decreased blood pressure (orthostatic hypotension). You may feel lightheaded or faint when you rise too quickly from a sitting or lying position.
- Falls. REXULTI may make you sleepy or dizzy, may cause a decrease in your blood pressure when changing position (orthostatic hypotension), and can slow your thinking and motor skills which may lead to falls that can cause fractures or other injuries.
- Seizures (convulsions).
- Problems controlling your body temperature so that you feel too warm. Do not become too hot or dehydrated during treatment with REXULTI. Do not exercise too much. In hot weather, stay inside in a cool place if possible. Stay out of the sun. Do not wear too much clothing or heavy clothing. Drink plenty of water.
- Difficulty swallowing that can cause food or liquid to get into your lungs.

(continued on next page)

IMPORTANT SAFETY INFORMATION for REXULTI® (brexpiprazole) (continued)

REXULTI may cause serious side effects, including: (continued)

• Sleepiness, drowsiness, feeling tired, difficulty thinking and doing normal activities. Do NOT drive a car, operate machinery, or do other dangerous activities until you know how REXULTI affects you. REXULTI may make you feel drowsy.

Before taking REXULTI, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had heart problems or a stroke
- have or had low or high blood pressure
- have or have had diabetes or high blood sugar or a family history of diabetes or high blood sugar. Your healthcare provider should check your blood sugar before you start REXULTI and during treatment with REXULTI.
- have or have had high levels of total cholesterol, LDL cholesterol, or triglycerides, or low levels of HDL cholesterol
- have or had seizures (convulsions)
- have or have had kidney or liver problems
- have or have had a low white blood cell count
- have or had a low white blood cell count
- are pregnant or plan to become pregnant. REXULTI may harm your unborn baby. Taking REXULTI during your third trimester of pregnancy may cause your baby to have abnormal muscle movements or withdrawal symptoms after birth.
 - o Tell your healthcare provider if you become pregnant or think you are pregnant during treatment with REXULTI.
 - If you become pregnant during treatment with REXULTI, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics. You can register by calling 1-866-961-2388 or visit http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/.
- are breastfeeding or plan to breastfeed. It is not known if REXULTI passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with REXULTI.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. REXULTI and other medicines may affect each other causing possible serious side effects. REXULTI may affect the way other medicines work, and other medicines may affect how REXULTI works. Your healthcare provider can tell you if it is safe to take REXULTI with your other medicines. Do not start or stop any medicines during treatment with REXULTI without first talking to your healthcare provider.

The most common side effects of REXULTI include weight gain and restlessness or feeling like you need to move (akathisia).

These are not all the possible side effects of REXULTI. For more information, ask your healthcare provider or pharmacist.

You are encouraged to report side effects of REXULTI (brexpiprazole). Please contact Otsuka America Pharmaceutical, Inc. at 1-800-438-9927 or FDA at 1-800-FDA-1088 (www.fda.gov/medwatch).

Please read **FULL PRESCRIBING INFORMATION**, including **BOXED WARNING**, and **MEDICATION GUIDE**.



